

ORIGINAL ARTICLE

THE USE OF IMMUNOENZYMATIC METHOD FOR DETECTION OF ANTIBODIES AGAINST ZOO NOTIC DISEASES IN CZECH SOLDIERS RETURNING FROM AFGHANISTAN

Petra Strakova^{1,2} ✉, Ivo Rudolf^{1,2}, Oto Pavlis³, Zdenek Hubalek^{1,2}

1 Institute of Vertebrate Biology, v.v.i., Academy of Sciences of the Czech Republic, Brno

2 Department of Experimental Biology, Faculty of Science, Masaryk University, Brno

3 Military Health Institute, Biological Defence Department, 56166 Těchonín

Received 6th May 2014.

Revised 20th May 2014.

Published 2nd June 2014.

Summary

In this work, we focused on detection of IgG antibodies in the blood sera from Czech soldiers returning from Afghanistan against selected zoonotic diseases by commercially available ELISA kits. Samples were tested for the presence of antibodies against *Brucella abortus*, *Coxiella burnetii*, *Leptospira* spp., complex of *Rickettsia conorii*, *Leishmania infantum*, hantaviruses and hepatitis E virus. Except for *L. infantum* (all persons were negative), we found following seroreactivity rate: 10 % in *B. abortus*, 11 % in *C. burnetii*, 20 % in *Leptospira* spp., 10 % in *R. conorii* and 4 % in both hepatitis E virus and hantaviruses.

Key words: ELISA method; zoonoses; soldiers; Afghanistan

INTRODUCTION

The zoonoses are infectious diseases that are transmitted to humans from animals. Their agents can be parasites, fungi, bacteria, viruses or prions [1]. We selected seven zoonotic agents that should be tested in soldiers returning from missions – *Brucella*, *Leishmania*, *Rickettsia*, *Leptospira*, *Coxiella*, hepatitis E virus, and hantavirus [2]. Afghanistan lies in several climate zones in the south-west Asia. Its surface is very ragged and hostile. Years of wars and civil disorders led to huge flood of soldiers, peacekeepers, humani-

tarian workers and journalists. Also the number of refugees and internally displaced persons increased, which, with the number of foreigners, enhanced the epidemiological risk, including zoonoses. The population in Afghanistan is affected by many infectious diseases – especially diarrhea, respiratory infections and measles [3]. Soldiers go to tropical or subtropical places with low hygiene standards and with the presence of many infectious diseases [4]. Despite of having clean water and food, clean place to stay and being vaccinated, the soldiers should take into account the risk of infection.

Bacteria of the genus *Brucella* are gram negative coccobacilli, causing a worldwide known disease called brucellosis. It affects primarily animals but can easily be transmitted to humans. The most dangerous species are *Brucella melitensis*, *B. abortus*, *B. suis*

✉ Academy of Sciences of the Czech Republic,
Institute of Vertebrate Biology, v.v.i.,
Kláštěrní 2, 691 42 Valtice, Czech republic
strakova.p@centrum.cz

and *B. canis* [5]. The reservoir of these bacteria can be cattle, goats, sheep, camels, pigs and some other animals. People can be infected by direct contact with animals, infected dust, contaminated milk or dairy products [6].

Coxiella belongs to gram negative and intracellular bacteria. *Coxiella burnetii* is the only member of this genus and it causes Q fever. It occurs worldwide and it is dangerous for both people and animals. People can get infected in the same ways as with *Brucella* and additionally by ticks [1].

Leptospira belongs to class *Spirochaetes* and it is gram negative. It causes disease called leptospirosis. Its hosts and reservoirs are rodents, insectivores, dogs, cattle, pigs, horses and many more. The most frequent way of infection is contact with contaminated water, soil or food [7].

Bacteria from the genus *Rickettsia* are small gram negative and intracellular organisms. They are agents of several diseases. They are associated with haematophagous arthropods (ticks, fleas, lice, chiggers) and some of them are reservoir and vectors of the disease. In the Middle East, only *Rickettsia conorii* occurs. It can cause Mediterranean fever and other fevers. In addition to high fever, the disease is characterized by rash and eschar [8].

Leishmania is a protozoon causing leishmaniasis that occurs endemically in tropical and subtropical regions. It is transmitted by phlebotomus. *Leishmania* species differ in disease symptoms and geographic distribution. Some of them cause cutaneous, mucocutaneous or visceral leishmaniasis.

Hantaviruses belong to the family *Bunyaviridae* and they occur in Asia, America, Africa and Europe.

They cause two main types of disease: haemorrhagic fever with renal syndrome (HFRS) and Hantavirus pulmonary syndrome (HPS). Their natural host and reservoir are rodents [1].

Hepatitis E virus is non-enveloped virus and it is the only member of the genus *Hepevirus*. The five genotypes are known and four of them occur in people [8]. The transmission is faecal-oral through contaminated water or food, but there is also a possibility of zoonotic transmission.

MATERIALS AND METHODS

Blood samples

For this study, the blood sera were collected from 70 asymptomatic persons staying in Afghanistan in 2011 after their arrival and returning to the Czech Republic and were supplied from Czech Army Serum Bank. The blood samples were centrifuged in the laboratory 3 to 6 hours after collection and the separated sera were stored at -20 °C.

ELISA assay

We used commercially available ELISA kits for determination of IgG antibodies (Table 1). These kits apply indirect ELISA method. Every test was performed according to manual provided by the producer.

After measurement of optical density (OD) by spectrophotometer, we calculated results by using formula provided in manual. These results are also interpreted according to a table found in each brochure, most often as a ratio of sample OD and the cut-off value.

Table 1. ELISA kits used in our testing.

Zoonotic agents	ELISA kit
<i>Brucella abortus</i>	Brucella-IgG, NovaTec Immunodiagnostica
<i>Coxiella burnetii</i>	Coxiella-IgG, DIAMedix, Delta Biologicals
<i>Leptospira</i> spp.	SERION ELISA classic IgG leptospira
<i>Rickettsia conorii</i>	Rickettsia-IgG, Fuller laboratories
<i>Leishmania infantum</i>	Leishmania-IgG, NovaTec Immunodiagnostica
Hantavirus Eurasia (Hantaan, Dobrava, Puumala)	Anti-Hanta virus pool 1 Eurasia, EUROIMMUN
Hepevirus	HEV IgG, DIA.PRO

RESULTS AND DISCUSSION

The results are given in Table 2.

Brucella abortus – out of 70 tested sera, 7 sera were positive (10 %) for the presence of IgG antibodies against *B. abortus* W 99. We can assume

that if there was an infection, it was mild or inapparent. Unfortunately, the producer doesn't provide an exact information about what antigen was used. In many cases, the immunotests mainly contain proteins of outer membrane. The most cases of human brucellosis are caused by *B. abortus* and *B. melitensis*. But because of genetic similarity

Table 2. Reactivity of the sera in ELISA (–, negative serum; +, positive serum).

No. of person	<i>Brucella abortus</i>	<i>Coxiella burnetii</i>	<i>Leptospira</i> sp.	<i>Rickettsia conorii</i>	<i>Leishmania infantum</i>	Hantavirus	Hepevirus
1	–	–	+	+	–	–	–
3	–	–	–	+	–	–	–
7	–	–	–	+	–	–	–
8	–	–	+	–	–	–	–
10	–	–	+	–	–	–	–
17	+	–	+	–	–	–	–
21	+	+	–	+	–	–	–
22	+	+	+	–	–	–	–
23	+	–	–	–	–	–	+
26	+	–	+	+	–	–	–
29	–	–	+	–	–	+	–
30	–	–	+	–	–	+	–
31	–	+	–	–	–	–	–
32	–	–	–	–	–	–	+
33	–	–	+	+	–	–	–
36	–	+	–	–	–	–	–
37	–	–	+	–	–	–	–
42	+	–	+	–	–	–	–
43	–	–	–	–	–	–	+
48	–	–	–	–	–	+	–
49	–	+	–	–	–	–	–
50	–	+	–	+	–	–	–
60	–	+	–	–	–	–	–
64	–	–	+	–	–	–	–
66	–	+	–	–	–	–	–
67	–	–	+	–	–	–	–
68	+	–	+	–	–	–	–
Σ (pos.)	7	8	14	7	0	3	3
% (pos.)	10	11.4	20	10	0	4.3	4.3

The sera no. 2, 4–6, 9, 11–16, 18–20, 24, 25, 27, 28, 34, 35, 38–41, 44–47, 51–59, 61–63, 65, 69 and 70 did not react with any of the seven antigens and are not given in the Table.

within the *Brucella* genus, we can assume that the different species will have the same or similar major outer membrane proteins (MOMP) [9]. The presence of brucellosis in the Middle East is endemic. Afghanistan belongs to countries with the highest incidence of brucellosis [10]. Some cases of brucellosis in the military units were described [11, 12, 13].

Coxiella burnetii – out of 70 tested sera, 8 sera were positive (11 %). Q fever is quite common in the Middle East and in Mediterranean. Because of many possible routes of transmission, the probability of infection in endemic regions is very high. Špliňo et al. [14] described an epidemic among Czech soldiers in Bosnia. Unfortunately, the literature about Q fever in Afghanistan is limited. Arsen'eva [15] tested animal sera. Among troops, authors mostly described cases from Iraq [16, 17, 18].

Leptospira spp. – out of 70 tested sera, 14 sera were positive (20 %). The producer states that this ELISA kit should detect genus-specific human antibodies against *Leptospira*. But the classification of the genus *Leptospira*, with so many serovars, is difficult so it is not possible to identify specific serovar in the positively reacting sera. Leptospirosis is dangerous for soldiers operating in swamps [19]. Pappas et al. [20] mentioned that there are no data from countries suffering from wars. Bryan et al. [21] tested 570 sera from soldiers in Pakistan and they found antibodies in 1-6 % of them. Mansour-Ghanaer et al. [22] found 74 cases of leptospirosis in Iran.

Rickettsia conorii – in this kit, the antigen was obtained from *Rickettsia conorii*. The producer states that this antigen reacts equally with *R. rickettsii*, *R. slovaca* and *R. africae*. We found 7 positive sera (10 %). It is known that there are two rickettsial diseases in Afghanistan – Siberian tick typhus and Mediterranean fever. Many authors described just the presence of different rickettsial species but only a few authors focused on the evidence of the disease [23, 24, 25, 26, 27].

Leishmania infantum – we found no positive sera. The reason of our result is that *L. infantum* causes visceral leishmaniasis which is found only sporadically in Afghanistan [28, 29]. Visceral leishmaniasis can be found in Iran or Pakistan [13]. On the other hand, Afghanistan is known for the presence of cutaneous leishmaniasis [30, 31, 32, 33].

Hantavirus – out of 70 tested sera, 3 sera were positive (4 %). This kit uses recombinant antigen of three hantaviruses (Hantaan, Dobrava and Puumala). The cases of HFRS were known from the past (Bosnia - Stuart et al. [34]; Croatia - Mulic and Ropac [35]). The literature about the Middle East is limited and sometimes the authors differ in their statements [4, 36].

Hepevirus – out of 70 tested sera, 3 sera were positive (4 %). The most cases of hepatitis E occur in south Asia [37], where the contaminated water is a usual source of infection. But without laboratory confirmation, it is very difficult to distinguish hepatitis E from other viral hepatitis [37]. About the situation in Afghanistan or in the soldiers, there are limited data [38, 39, 40]. Carmoi et al. [41] tested Afghan population and they found high prevalence of antibodies – 28.4 %. Other authors focused on neighboring countries (Pakistan, Iran), so we can assume that hepatitis E might be endemic [42, 43, 44, 45].

For diagnosis of some zoonotic diseases, ELISA method is not available or ELISA method is not used at all (cutaneous leishmaniasis).

CONCLUSIONS

In this study, we focused on detection of specific IgG antibodies against selected zoonotic agents with which the Czech soldiers can come across during their mission in Afghanistan. We tested blood samples from 70 returning soldiers by commercially available ELISA kits. Seven sera (10 %) reacted with antigen from *Brucella abortus*, eight sera (13 %) reacted with antigen from *Coxiella burnetii*, fourteen sera (20 %) reacted with antigen from *Leptospira* spp., seven sera (10 %) reacted with antigen from *Rickettsia* spp., no serum reacted with antigen from *Leishmania infantum*, three sera (4 %) reacted with antigen from Hantavirus and three sera (4 %) reacted with antigen from hepatitis E virus.

After their arrival, all soldiers were medically examined and we know that all soldiers, we tested, were found healthy. We have no information about their position or health condition during their mission and about their previous missions. We can only assume that they came into contact with selected zoonotic agents during their last mission in Afghanistan.

In literature we did not find enough data about the epidemiological situation in Afghanistan (including zoonoses), probably due to the war conflict. According to our results we can assume that in Afghanistan there is a chance for possible infection of soldiers or travelers with some zoonotic agents. In many cases these organisms cause only fever or the infection could be asymptomatic and the proper laboratory diagnosis should rely on the combination of cultivation, microscopy, molecular techniques and not only on serological data.

ACKNOWLEDGEMENTS

This study was partially funded by the EU grant FP7-261504 EDENext (<http://www.edenext.eu>). The publication is catalogued by the EDENext Steering Committee as EDENext229. The contents of this paper are the sole responsibility of the authors and do not necessarily reflect the views of the European Commission.

REFERENCES

- Hubálek, Z.; Rudolf, I. Microbial Zoonoses and Saprozooses. Springer, Dordrecht-Heidelberg-London-New York **2011**, 457 pp.
- Smetana, J.; Smetana, R.; Vacková, M. Protiepidemické zabezpečení vojáků Armády České republiky před výjezdem a po návratu ze zahraniční mise. *Voj. zdravot. Listy*. **2004**, 73, 41-44.
- Assefa, F.; Jabarkhil, M.; Salama, P. Malnutrition and mortality in Kohistan District, Afghanistan. *J. Am. Med. Assoc.* **2001**, 286, 2723-2728.
- Wallace, M.R.; Hale, B.R.; Utz, G.C.; Olson, P.E.; Earhart, K.C.; Thornton, S.A.; Hyams, K.C. Endemic Infectious Diseases of Afghanistan. *Clin. Infect. Dis.* **2002**, 34, 171-207.
- Maloney, G.E.; Mothershead, J.L.; Talavera, F.; Kulkarni, R.; Halamka, J.D.; Darling, R.G. CBRNE-Brucellosis **2009**. Dostupné na: <http://emedicine.medscape.com/article/830118-overview>, staženo 17.2.2013.
- Memish, Z.A.; Almuneeef, M.; Mah, M.W.; Qassem, L.A.; Osoba, A.O. Comparison of the *Brucella* standard agglutination test with the ELISA IgG and IgM in patients with *Brucella* bacteremia. *Diagn. Microbiol. Infect. Dis.* **2002**, 44, 129-132.
- Stoddard, R.; Shadomy, S.V. Leptospirosis 2003. In: Yellow book. Dostupné na: <http://wwwnc.cdc.gov/travel/yellowbook/2012/chapter-3-infectious-diseases-related-to-travel/leptospirosis.htm>
- Kasper, D.L.; Fauci, A.S. Harrison's Infectious Diseases. The McGraw-Hill Companies, New York **2010**, 1294 pp.
- Hunter, S.B.; Bibb, W.F.; Shih, C.N.; Kaufmann, A.F.; Mitchell, J.R.; McKinney, R.M. Enzyme-linked immunosorbent assay with major outer membrane proteins of *Brucella melitensis* to measure immune response to *Brucella* species. *J. Clin. Microbiol.* **1986**, 24, 566-572.
- Pappas, G.; Papadimitriou, P.; Akritidis, N.; Christou, L.; Tsianos, E.V. The new global map of human brucellosis. *Lancet*. **2006**, 6, 91-99.
- Bechtol, D.; Carpenter, L.R.; Mosites, E.; Smalley, D.; Dunn, J.R. *Brucella melitensis* infection following military duty in Iraq. *Zoonos. Publ. Hlth.* **2011**, 58, 489-492.
- Andrews, R. Brucellosis in a soldier who recently returned from Iraq, July 2004. *Med. Surveill. Monthly Rep.* 2004, 10, 30. Quoted by Aronson et al. **2006**.
- Aronson, N.E.; Sanders, J.W.; Moran, K.A. In harm's way: infections in deployed American military forces. *Clin. Inf. Dis.* **2006**, 43, 1045-1051.
- Šplíňo, M.; Beran, J.; Chlíbek, R. Q fever outbreak during the Czech Army deployment in Bosnia. *Mil. Med.* **2003**, 168, 840.
- Arsen'eva, L.P. Natural focus of zoonoses in Afghanistan (in Russian). *Med. Parazitol. (Mosk)*. **1982**, 51, 54-59.
- Anderson, A.D.; Smoak, B.; Shuping, E.; Ockenhouse, C.; Petrucci, B. Q fever and the US Military. *Emerg. Infect. Dis.* **2005**, 11, 1320-1322.
- Faix, D.J.; Harrison, D.J.; Riddle, M.S.; Vaughn, A.F.; Yingst, S.L.; Earhart, K.; Thibault, G. Outbreak of Q fever among US military in western Iraq, June-July 2005. *Clin. Infect. Dis.* **2008**, 46, 65-68.
- Anderson, A.D.; Baker, T.R.; Littrell, A.C.; Mott, R.L.; Niebuhr, D.W.; Smoak, B.L.; Seroepidemiologic survey for *Coxiella burnetii* among hospitalized US troops deployed to Iraq. *Zoonos. Publ. Hlth.* **2011**, 58, 276-283.
- Corwin, A.; Ryan, A.; Bloys, W. A waterborne outbreak of leptospirosis among United States military personnel in Okinawa, Japan. *Int. J. Epidemiol.* **1990**, 19, 743.
- Pappas, G.; Papadimitriou, P.; Siozopoulou, V.; Christou, L.; Akritidis, N. The globalization of leptospirosis: worldwide incidence trends. *Int. J. Inf. Dis.* **2008**, 12, 351-357.

21. Bryan, J.P.; Mohammed, I.; Ksiazek, T.G.; Aftab, A.; Duncan, J.F.; Bashir, A.; Krieg, R.E.; Mohammed, R.; Leduc, J.W.; Shahid, N.; Shuaib, Q.; Malik, I.A.; Legters, L.J. Prevalence of sand fly fever, West Nile, Crimean-Congo hemorrhagic fever, and leptospirosis antibodies in Pakistan military personnel. *Mil. Med.* **1996**, *161*, 149-153.
22. Mansour-Ghanaei, F.; Sarshad, A.; Fallah, M.S.; Pourhabibi, A.; Pourhabibi, K.; Yousefi-Mashhoo, M. Leptospirosis in Guilan, a northern province of Iran: assessment of the clinical presentation of 74 cases. *Med. Sci. Monit.* **2005**, *11*, 219-223.
23. Raoult, D.; Roux, V. Rickettsioses as paradigms of new or emerging infectious diseases. *Clin. Microbiol.* **1997**, *10*, 694-719.
24. Robertson, R.G.; Wisseman, C.L. Tick-borne rickettsiae of the spotted fever group in West Pakistan. II. Serological classification of isolates from West Pakistan and Thailand: evidence for two new species. *Am. J. Epidemiol.* **1973**, *97*, 55-64.
25. Fournier, P.E.; Durand, J.P.; Rolain, J.M.; Camicas, J.L.; Tolou, H.; Raoult, D. Detection of Astrakhan fever rickettsia from ticks in Kosovo. *Ann. N.Y. Acad. Sci.* **2003**, *990*, 158-161.
26. Kelly, D.J.; Richards, A.L.; Temenak, J.; Strickman, D.; Dasch, G.A. The past and present threat of rickettsial diseases to military medicine and international public health. *Clin. Infect. Dis.* **2002**, *34*, 145-169.
27. Walker, D.H. Rickettsial diseases in travelers. *Travel Med. Infect. Dis.* **2003**, *1*, 35-40.
28. Leslie, T.; Saleheen, S.; Sami, M.; Mayan, I.; Mahboob, N.; Fiebert, K. Visceral leishmaniasis in Afghanistan. *Can. Med. Assoc. J.* **2006**, *175*, 245.
29. Myles, O.; Wortmann, G.W.; Cummings, J.F.; Barthel, V.; Patel, S.; Crum-Cianflone, N.F.; Negin, N.S.; Weina, P.J.; Ockenhouse, C.F.; Joyce, D.J.; Magill, A.J.; Aronson, N.E.; Gasser, R.S. Visceral leishmaniasis: clinical observations in 4 US army soldiers deployed to Afghanistan or Iraq, 2002-2004. *Arch. Intern. Med.* **2007**, *167*, 1899-1901.
30. Reithinger, R.; Mohsen, M.; Aadil, K.; Sidiqi, M.; Erasmus, P.; Coleman, P.G. Anthroponotic cutaneous leishmaniasis, Kabul, Afghanistan. *Emerg. Infect. Dis.* **2007**, *9*, 727-729.
31. Faulde, M.K.; Heyl, G.; Amirih, M.L. Zoonotic cutaneous leishmaniasis, Afghanistan. *Emerg. Infect. Dis.* **2006**, *12*, 1623-1624.
32. Faulde, M.; Schrader, J.; Heyl, G.; Amirih, M. Differences in transmission seasons as an epidemiological tool for characterization of anthroponotic and zoonotic cutaneous leishmaniasis in northern Afghanistan. *Acta Trop.* **2008**, *105*, 131-138.
33. Van Thiel, P.; Leenstra, T.; Kager, P.A.; de Vries, H.J.; van Vugt, M.; van der Meide, W.F.; Bart, A.; Zeegelaar, J.E.; van der Sluis, A.; Schallig, H.D.F.H.; van Gool, T.; de Vries, P.J. Miltefosine treatment of Leishmania major infection: an observational study involving dutch military personnel returning from northern Afghanistan. *Clin. Inf. Dis.* **2010**, *50*, 80-83.
34. Stuart, L.M.; Rice, P.S.; Lloyd, G.; Beale, R. A soldier in respiratory distress. *Lancet.* **1996**, *346*, 30.
35. Mulic, R.; Ropac, D. Epidemiologic characteristic and military implications of hemorrhagic fever with renal syndrome in Croatia. *Croat. Med.* **2002**, *43*, 581.
36. Lee, H.W. Hemorrhagic fever with renal syndrome in Korea. *Clin. Infect. Dis.* **1989**, *11*, 846-876.
37. Labrique, A.B.; Thomas, D.L.; Stoszek, S.K.; Nelson, K.E. Hepatitis E: an emerging infectious disease. *Epidemiol. Rev.* **1999**, *21*, 162-179.
38. Eick, A.; Ticehurst, J.; Tobler, S.; Nevin, R.; Lindler, L.; Hu, Z.; Macintosh, V.; Jarman, R.G.; Gibbons, R.V.; Myint, K.S.A.; Gaydos J. Hepatitis E seroprevalence and seroconversion among US military service members deployed to Afghanistan. *J. Infect. Dis.* **2010**, *202*, 1302-1308.
39. Labrique, A.B.; Zaman, K.; Hossain, Z.; Saha, P.; Yunus, M.; Hossain, A.; Ticehurst, J.R.; Nelson, K.E. Epidemiology and risk factors of incident hepatitis E virus infections in rural Bangladesh. *Am. J. Epidemiol.* **2010**, *172*, 952-961.
40. Myint, K.S.; Endy, T.P.; Shrestha, M.P. Hepatitis E antibody kinetics in Nepalese patients. *Trans. R. Soc. Trop. Med. Hyg.* **2006**, *100*, 938-941.
41. Carmoi, T.; Safiullah, S.; Nicand, E. Risk of enterically transmitted hepatitis A, hepatitis E and Plasmodium falciparum malaria in Afghanistan. *Clin. Infect. Dis.* **2009**, *48*, 1800.
42. Alavian, S.M.; Fallahian, F.; Lankarani, K.B. Epidemiology of hepatitis E in Iran and Pakistan. *Hepat. Mon.* **2009**, *9*, 60-65.
43. Ataei, B.; Nokhodian, Z.; Javadi, A.A.; Kassaian, N.; Shoaie, P.; Farajzadegan, Z.; Adibi, P. Hepatitis E virus in Isfahan province: a population-based study. *Int. J. Inf. Dis.* **2008**, *13*, 67-71.
44. Malik, I.A.; Tariq, W.U.Z. The prevalence and pattern of viral hepatitis in Pakistan. *J. College Physicians Surgeons Pakist.* **1995**, *5*, 2.
45. Bryan, J.P.; Iqbal, M.; Tsarev, S.; Malik, I.A.; Duncan, F.; Ahmed, A.; Khan, A.; Rafiqui, A.R.; Purcell, R.H.; Legters, L.J. Epidemic of hepatitis E in a military unit in Abbottabad, Pakistan. *Am. J. Trop. Med. Hyg.* **2002**, *67*, 662-668.